

Single parameter quasi-particle model for QGP

Vishnu M. Bannur

Department of Physics,

University of Calicut, Kerala-673 635, India.

February 2, 2008

Abstract

We discuss a new single parameter quasi-particle model and study the thermodynamics of (2+1)-flavor quark gluon plasma (QGP). Our model with a single parameter explains remarkably well the lattice simulation results of Fodor *et. al.* [1].

PACS Nos : 12.38.Mh, 12.38.Gc, 05.70.Ce, 52.25.Kn

Keywords : Equation of state, quark gluon plasma, quasi-particle plasma

1 Introduction :

Quasi-particle model (qQGP) of quark gluon plasma (QGP) was first proposed by Peshier *et. al.* [2] to explain the non-ideal equation of state (EoS), observed in lattice gauge theory simulations (LGT) [3]. At finite temperature, instead of real quarks and gluons *with* QCD (quantumchromodynamics) *interactions* we may as well consider the system to be made up of *non-interacting* quasi-particles with thermal masses, quasi-quarks and quasi-gluons, and study the thermodynamics. Quasi-particles are quanta of plasma collective modes excited by quarks and gluons through QCD interactions. Initial quasi-particle model was found to be thermodynamically inconsistent [4] and also not able to fit the more recent LGT results [5]. Gorenstein and Yang reformulated the statistical mechanics (SM) to solve the inconsistency, but end up with an extra undetermined, temperature dependent terms in the expressions for pressure and energy density, which need to be phenomenologically chosen. However, as we discuss here [6], above reformulation of SM is not needed and standard SM may be applied to qQGP without any extra phenomenological terms. There is no TD inconsistency in our new qQGP model. Peshier's model with reformulated SM by Gorenstein and Yang has been studied by various groups [7, 8, 9, 10, 11] with different expressions for thermal masses, effective degrees of freedom, so on. Note that all of above works are based on the reformulation of SM of Gorenstein and Yang, which in fact based on mathematical identities involving derivatives with respect to temperature and chemical potentials, used to redefine average energy density (ε) and number density (n) respectively. As we have shown recently in Ref. [6], we may skip this TD inconsistency problem and instead use the original definition of ε and n , and making use of TD relations we may get all TD quantities. As a specific example, here we discuss (2+1)-flavor QGP, studied by Fodor *et. al.* [1] using LGT.

Of course, there are other models like HTL (hard thermal loop) [12], recent FMR (fundamental modular region) gas [13] etc. based on QCD perturbative and non-perturbative calculations, but fails to fit LGT results near to the transition temperature T_c . At the same time phenomenological models of QGP, based on plasma theory with QCD inputs like SCQGP (strongly coupled quark gluon plasma) [14], our present qQGP model seems to fit

remarkably well the LGT results with minimum number of parameters. All other qQGP models, field theoretical models [15], Ploykov loop models [16] also fit LGT results, but by adjusting 3 or more parameters. Of course, we know that near $T = T_c$, region of phase transition or cross over, it is a low energy phenomena and hence QCD can not be solved by analytical methods like perturbation theory because the coupling constant α_s is not small enough. Probably we need to formulate phenomenological models, just like in the case of hadron spectroscopy, to study TD of QGP near T_C .

2 Phenomenological Model with $\mu = 0$:

QGP at thermodynamic equilibrium consists of interacting quarks and gluons which exhibits collective behaviour. Our basic assumption is that this system may be replaced by a system of non-interacting quasi-particles with quantum numbers of quarks and gluons. These quasi-particles have additional thermal masses which are equal to plasma frequencies. Here we differ from other qQGP models where, for example, the thermal mass was taken to be $\sqrt{3/2}$ times the plasma frequency. A general expression for thermal mass or polarization tensor is very complicated expressions which is a function of momentum and frequency. Only at high momentum limit it approaches a simpler form which on further approximations reduces to above form. In view of such a drastic approximation and since we use phenomenological model one may as well take $m_{th} \approx \omega_p$. In fact with this relation, we get better result than with $m_{th} \approx \sqrt{3/2}\omega_p$. Further important point is that the above dispersion relation is obtained using perturbation methods with temperature dependent density distribution function appropriate to ideal system. Then one formulates TD of a non-ideal system. In principle, this must be carried out in a self-consistent manner as discussed in Ref. [17], where, for example, density expression is an integral equation since ω_p depends on density. So we need to solve an integral equation self-consistently to get the the density. Here we avoid all above complications and as a phenomenological input, we assume that $m_{th} = \omega_p$.

Following the standard procedure of statistical mechanics [18], the grand partition func-

tion is given by,

$$Q_G = \sum_{s,r} e^{-\beta E_r - \alpha N_s} , \quad (1)$$

where the sum is over energy states E_r and particle number states N_s . α and β are defined as $\alpha \equiv -\mu/T$ and $\beta \equiv 1/T$. Next on further simplifications and taking thermodynamic limit, we get,

$$q \equiv \ln Q_G = \mp \sum_{k=0}^{\infty} \ln(1 \mp z e^{-\beta \epsilon_k}) , \quad (2)$$

where q is called q-potential and $\mp$ for bosons and fermions. $z \equiv e^{\mu/T} = e^{-\alpha}$ is called fugacity. ϵ_k is the single particle energy, given by,

$$\epsilon_k = \sqrt{k^2 + m^2} ,$$

where k is momentum and m is the total mass which contains both the rest mass and thermal mass (m_{th}) of particles. m_{th} may depend on temperature T and chemical potential μ depending on QGP system. All other qQGP models starts from q-potential to get pressure with correction term to avoid TD inconsistency and proceed to evaluate other TD quantities. However, we avoid this trap of inconsistency problem by adopting a different procedure. We know that, for grand canonical ensemble, TD quantities like energy density and number density may be obtained by taking ensemble averages. That is, the average energy U is given by,

$$U \equiv \langle E_r \rangle = \frac{\sum_{s,r} E_r e^{-\beta E_r - \alpha N_s}}{Q_G} = -\frac{\partial}{\partial \beta} \ln Q_G = \sum_k \frac{z \epsilon_k e^{-\beta \epsilon_k}}{1 \mp z e^{-\beta \epsilon_k}} . \quad (3)$$

Note that the partial differentiation with respect to β above is just a mathematical trick to express U in terms of sum over single particle energy levels, ϵ_k , making use of Eq. (2). While differentiating, indirect dependence of $\beta = 1/T$ in the fugacity, z , and thermal mass, $m_{th}(T, \mu)$, must be ignored by definition. Otherwise, we will not get back $\langle E_r \rangle$. Now, on taking continuum limit and after some algebra, we get,

$$\epsilon = \frac{g_f T^4}{2 \pi^2} \sum_{l=1}^{\infty} (\pm 1)^{l-1} z^l \frac{1}{l^4} \left[\left(\frac{m l}{T} \right)^3 K_1 \left(\frac{m l}{T} \right) + 3 \left(\frac{m l}{T} \right)^2 K_2 \left(\frac{m l}{T} \right) \right] , \quad (4)$$

where g_f is the degeneracy and equal to $g_g \equiv 16$ for gluons and equal to $12 n_f$ for quarks. n_f is the number of flavors with same mass. K_1 and K_2 are modified Bessel functions of order 1 and 2 respectively.

Let us now consider our main topic, (2+1)-flavor system, studied by Fodor *et. al.* [1], using our model. It is a QGP with two light (u) and one heavy (s) quarks along with gluons. Let us first consider the case with zero chemical potential and take $z = 1$. Hence we get the energy density, expressed in terms of $e(T) \equiv \varepsilon/\varepsilon_s$, for the quark gluon plasma of quasi-partons is

$$\begin{aligned}
e(T) = & \frac{15}{\pi^4} \frac{1}{(g_f + \frac{21}{2} n_f^{eff})} \sum_{l=1}^{\infty} \frac{1}{l^4} \left(g_f \left[\left(\frac{m_g l}{T} \right)^3 K_1 \left(\frac{m_g l}{T} \right) + 3 \left(\frac{m_g l}{T} \right)^2 K_2 \left(\frac{m_g l}{T} \right) \right] \right. \\
& + 12 n_u^{eff} (-1)^{l-1} \left[\left(\frac{m_u l}{T} \right)^3 K_1 \left(\frac{m_u l}{T} \right) + 3 \left(\frac{m_u l}{T} \right)^2 K_2 \left(\frac{m_u l}{T} \right) \right] \\
& \left. + 12 n_s^{eff} (-1)^{l-1} \left[\left(\frac{m_s l}{T} \right)^3 K_1 \left(\frac{m_s l}{T} \right) + 3 \left(\frac{m_s l}{T} \right)^2 K_2 \left(\frac{m_s l}{T} \right) \right] \right) , \tag{5}
\end{aligned}$$

where ε_s is the Stefan-Boltzman gas limit of QGP, which may be obtained by taking high temperature limits of Eq. (4) for gluons and quarks separately and adding them. m_g is the temperature dependent gluon mass (m_{th}), which is equal to the plasma frequency, i.e, $m_g^2 = \omega_p^2 = \frac{g^2 T^2}{18} (2N_c + n_f)$. All quarks have both the thermal mass as well as the rest mass and hence, the total mass may be written as

$$m_q^2 = m_{q0}^2 + \sqrt{2} m_{q0} m_{th} + m_{th}^2 , \tag{6}$$

following the idea used in other qQGP models for the system with finite quark masses. Only the difference is that our m_{th} is equal to the plasma frequency due quarks alone. That is, $m_{th}^2 = \omega_p^2 = \frac{g^2 T^2}{18} n_f$. m_{q0} is the rest mass of up or strange quark. g^2 in thermal masses is related to the two-loop order running coupling constant, given by,

$$\alpha_s(T) = \frac{6\pi}{(33 - 2n_f) \ln(T/\Lambda_T)} \left(1 - \frac{3(153 - 19n_f) \ln(2 \ln(T/\Lambda_T))}{(33 - 2n_f)^2 \ln(T/\Lambda_T)} \right) , \tag{7}$$

where Λ_T is a parameter related to QCD scale parameter. This choice of $\alpha_s(T)$ is motivated from lattice simulations. Using thermal masses with above α_s , we can evaluate the $e(T)$ from

Eq. (4). Note that the only temperature dependence in $e(T)$ comes from $\alpha_s(T)$, which has the same form as that of lattice simulations [19] with Λ_T as a free parameter. n_u^{eff} and n_s^{eff} are effective number of quarks which differ from 2 and 1, respectively, because of finite masses of up and strange quarks. However, handling of finite rest masses of quarks are different in LGT studies of Fodor *et. al.* and Bielefeld group [19]. Bielefeld group carried out the simulation with the ratio m_{q0}/T equal to constant and it is straight forward to calculate n_q^{eff} , where as Fodor *et. al.* used constant m_{q0} and it is not clear how to calculate n_q^{eff} . Hence as done in Ref. [20] we take in our calculations $n_u^{eff} = 2$, $n_s^{eff} = 0.5$ and $n_f^{eff} = 2.5$. Similar normalization need to be made to fit the LGT results in other models also, either multiplying LGT data with a factor 1.1 [9] or model's data with .9 [11], or sometime using massive gluon [21] so on. From ε , we may obtain pressure P by using a TD relation $\varepsilon = T \frac{\partial P}{\partial T} - P$ for $\mu = 0$ system and we get

$$\frac{P}{T} = \frac{P_0}{T_0} + \int_{T_0}^T dT \frac{\varepsilon(T)}{T^2} , \quad (8)$$

where P_0 and T_0 are pressure and temperature at some reference points. Results are presented in Fig. 1 along with LGT results. Note that earlier this phenomenological new qQGP model with a single system dependent adjustable parameter explained very well the LGT results of Bielefeld group [19] on QGP system with massless quarks as discussed in Ref. [6].

Another TD quantity which may be obtained from ensemble averaging is the number density, given by,

$$\langle N \rangle = \frac{\sum_{s,r} N_s e^{-\beta E_r - \alpha N_s}}{Q_G} = -\frac{\partial}{\partial \alpha} \ln Q_G = z \frac{\partial}{\partial z} \ln Q_G = \sum_k \frac{z e^{-\beta \epsilon_k}}{1 \mp z e^{-\beta \epsilon_k}} , \quad (9)$$

and using the definition $n \equiv \langle N \rangle / V$, where V is the volume.

3 Model with finite μ :

Let us next consider (2+1)-flavor system with finite μ and LGT results are available for quark density n_q or baryon density (n_B) and difference in pressure from $\mu = 0$ case ($\Delta P \equiv P(T, \mu) - P(T, \mu = 0)$). Here μ is the quark chemical potential which is one third of the baryon chemical potential. The present LGT results are with μ_B coming from up quarks

only and hence we need to consider only *up* quark density. Using the expression for the number density, Eq. (9), which on continuum limit and after some algebra, reduces to

$$\frac{n_q}{T^3} = \frac{12}{\pi^2} \sum_{l=1}^{\infty} (-1)^{l-1} \frac{1}{l^3} \left[\left(\frac{m_q l}{T} \right)^2 K_2 \left(\frac{m_q l}{T} \right) \sinh \left(\frac{\mu l}{T} \right) \right] . \quad (10)$$

Now we modify earlier $m_{th}^2(T)$ to $m_{th}^2(T, \mu)$ as

$$m_{th}^2(T, \mu) = \frac{g^2 T^2}{18} n_f \left(1 + \frac{\mu}{\pi^2 T^2} \right) , \quad (11)$$

inspired by QCD perturbative calculations [2]. In our case $n_f = 2$, mainly due to *up* quarks, and g^2 is related to two-loop order running coupling constant, discussed earlier, but need to be modified to take account of finite μ . Following the work of Schneider [22] and Letessier, Rafelski [21], now we change T/Λ_T in Eq. (7) as

$$\frac{T}{\Lambda_T} \sqrt{1 + a \frac{\mu^2}{T^2}} , \quad (12)$$

where a is a parameter which is equal to $(1.91/2.91)^2$ in the calculation of Schneider and $1/\pi^2$ in a phenomenological model of Letessier and Rafelski. In our model Schneider's $\alpha_s(T, \mu)$ works well.

From n_q , we may obtain other thermodynamic quantities like,

$$\Delta P \equiv P(T, \mu) - P(T, 0) = \int_0^\mu n_q d\mu , \quad (13)$$

and so on.

4 Results :

In Fig. 1, we plotted P/T^4 Vs T/T_c for (2+1)-flavor QGP with $\mu = 0$ and compared with LGT data. We took effective number of flavors as $n_u^{eff} = 2$, $n_s^{eff} = .5$ and $n_f^{eff} = 2.5$ because of finite quark masses. Quark rest masses are $m_{u0} = 65 \text{ MeV}$ and $m_{s0} = 165 \text{ MeV}$, values used in LGT simulations. Only the parameter $t_0 \equiv \Lambda_T/T_c$ is adjusted to get the best fit to LGT data on pressure and is equal to 0.4. In Fig. 2, the baryon density (n_B/T^3) is plotted as a function of T/T_c for different values of baryon chemical potentials and compared with LGT results. Surprisingly good fit is obtained without any new parameters. In Fig. 3, $\Delta P/T^4$ is plotted and again very good fit to LGT.

5 Conclusions :

Using our new formulation of qQGP phenomenological model, we were able to explain LGT results on (2+1)-flavor QGP with just a single parameter which may be related to QCD scale parameter. Our formalism is thermodynamically consistent and no need of reformulation of SM with extra temperature dependent terms in pressure and energy density so on. Earlier, using this model, we explained successfully the LGT results of Bielefeld group on QGP with massless quarks [6]. Hence a simple model of qQGP with a single parameter, related to QCD scale parameter, explains all existing LGT results and may explain future results in LGT and relativistic heavy ion collisions.

Of course, there are many other models which all claims to fit the LGT results, but all of them involve more than one parameter. Models with minimum number of parameters, which fits LGT results well, are SCQGP [14] and liquid model [20, 21], both with two parameters. All other models involve more than two parameters. Models based on QCD perturbative and non-perturbative calculations [12, 13] fails to fit the LGT results. Hence, it seems, phenomenological models based on the properties of plasma with QCD inputs explains well the LGT results.

References

- [1] Z. Fodor, S. D. Katz and K. K. Szabo, Phys. Lett. **B568**, 73 (2003).
- [2] A. Peshier, B. Kampfer, O. P. Pavlenko and G. Soff, Phys. Lett. **B337**, 235 (1994).
- [3] D. H. Rischke, M. I. Gorenstein, A. Schafer, H. Stocker and W. Greiner Phys. Lett. **B278**, 19 (1992); Z. Phys. **C56**, 325 (1992).
- [4] M. I. Gorenstein and S. N. Yang, Phys. Rev. **D52**, 5206 (1995).
- [5] V. M. Bannur, Eur. Phys. J. **C11**, 169 (1999).
- [6] V. M. Bannur, hep-ph/0508069.

- [7] A. Peshier, B. Kampfer, O. P. Pavlenko and G. Soff, Phys. Rev. **D54**, 2399 (1996); A. Peshier, B. Kampfer and G. Soff, Phys. Rev. **C61**, 045203 (2000).
- [8] P. Levai and U. Heinz, Phys. Rev. **C57**, 1879 (1998).
- [9] R. A. Schneider and W. Weise, Phys. Rev. **C64**, 055201 (2001).
- [10] K. K. Szabo and A. I. Toth, JHEP **0306**, 008 (2003).
- [11] Y. B. Ivanov, V. V. Skokov and V. D. Toneev, Phys. Rev. **D71**, 014005 (2005).
- [12] J. P. Blaizot, E. Iancu and A. Rebhan, Phys. Rev. Lett. **83**, 2906 (1999).
- [13] D. Zwanziger, Phys. Rev. Lett. **94**, 182301 (2005).
- [14] V. M. Bannur, hep-ph/0504072.
- [15] C. Ratti, M. A. Thaler and W. Weise, Phys. Rev. **D73**, 014019 (2006).
- [16] D. H. Rischke, Prog. Part. Nucl. Phys. **52**, 197 (2004).
- [17] V. M. Bannur, hep-ph/0602222.
- [18] R. K. Pathria, *Statistical Mechanics*, Butterworth Heinemann, Oxford (1997).
- [19] G. Boyd, J. Engels, F. Karsch, E. Laermann, C. Legeland, M. Lutgemeier and B. Petersson, Phys. Rev. Lett. **75**, 4169 (1995); Nucl. Phys. **B469**, 419 (1996); F. Karsch, Nucl. Phys. A, **698**, 199 (2002); E. Laermann and O. Philipsen, Ann. Rev. Nucl. Part. Sci. **53**, 163 (2003); F. Karsch, E. Laermann and A. Peikert Phys. Lett, **B478**, 447 (2000).
- [20] S. Hamieh, J. Letessier and J. Rafelski, Phys. Rev. **C62**, 064901 (2000).
- [21] J. Letessier and J. Rafelski, Phys. Rev. **C67**, 031902 (2003).
- [22] R. A. Schneider, hep-ph/0303104.

Figure 1: Plots of P/T^4 as a function of T/T_c from our model and lattice results (symbols) [1] for (2+1)-flavor QGP.

Figure 2: Plots of n_B/T^3 as a function of T/T_c from our model of (2+1)-flavor QGP and also with lattice data (symbols) [1] for $\mu_b = 100, 210, 330, 410$ and 530 MeV from bottom to top.

Figure 3: Plots of $\Delta P/T^4$ as a function of T/T_c from our model of (2+1)-flavor QGP and also with lattice data (symbols) [1] for $\mu_b = 100, 210, 330, 410$ and 530 MeV from bottom to top.




Fig. 1




Fig. 2




Fig. 3